

TESTING THE UNIVERSAL STRUCTURED JET MODELS OF GAMMA-RAY BURSTS BY BATSE OBSERVATIONS

XIAOHONG CUI^{1,4}, ENWEI LIANG^{1,2,3}, AND RUIJING LU^{1,2,4}

¹National Astronomical Observatories/Yunnan Observatory, Chinese Academy of Sciences, Kunming 650011, China

²Physics Department, Guangxi University, Nanning 530004, China

³Physics Department, University of Nevada, Las Vegas, NV89154, USA; Email: lew@physics.unlv.edu

⁴The Graduate School of the Chinese Academy of Sciences, Beijing, China

Draft version February 2, 2008

ABSTRACT

Assuming that the observed gamma-ray burst (GRB) rate as a function of redshift is proportional to a corrected star formation rate, we derive the empirical distribution of the viewing angles of long BATSE GRBs, $P^{\text{em}}(\theta)$, and the distribution of these bursts in the plane of θ against redshift, $P^{\text{em}}(\theta, z)$, by using a tight correlation between collimation-corrected gamma-ray energy (E_γ) and the peak energy of νF_ν spectrum measured in the rest frame (E'_p). Our results show that $P^{\text{em}}(\theta)$ is well fitted by a log-normal distribution centering at $\log \theta/\text{rad} = -0.76$ with a width of $\sigma_{\log \theta} = 0.57$. We test different universal structured jet models by comparing model predictions, $P^{\text{th}}(\theta)$ and $P^{\text{th}}(\theta, z)$, with our empirical results. To make the comparisons reasonable, an “effective” threshold, which corresponds to the sample selection criteria of the long GRB sample, is used. We find that (1) $P^{\text{th}}(\theta)$ predicted by a power-law jet model is well consistent with $P^{\text{em}}(\theta)$, but $P^{\text{th}}(\theta, z)$ predicted by this model is significantly different from $P^{\text{em}}(\theta, z)$; (2) $P^{\text{th}}(\theta, z)$ predicted by a single Gaussian jet model is more consistent with $P^{\text{em}}(\theta, z)$ than that predicted by the power-law jet model, but $P^{\text{th}}(\theta)$ predicted by this model rapidly drops at $\theta > 0.3$ rad, which greatly deviates $P^{\text{em}}(\theta)$; and (3) both $P^{\text{th}}(\theta)$ and $P^{\text{th}}(\theta, z)$ predicted by a two-Gaussian jet model are roughly consistent with our empirical results. A brief discussion shows that cosmological effect on the $E_\gamma - E'_p$ relation does not significantly affect our results, but sample selection effects on this relationship might significantly influence our results.

Subject headings: gamma rays: bursts—gamma rays: observations—methods: statistical

1. INTRODUCTION

The phenomenon of gamma-ray burst (GRB) is still of a great mystery, although significant progress has been made in the recent decade (see reviews by Fishman & Meegan 1995; Piran 1999; van Paradijs, Kouveliotou, & Wijers 2000; Cheng & Lu 2001; Mészáros 2002; Zhang & Mészáros 2004; Piran 2005). It is widely believed today that the central engines of GRBs power conical ejecta (jet) to produce the observed GRBs and their afterglows. Sharp breaks and/or quick decays of afterglow light curves are regarded as evidence of jets (e.g., Rhoads 1999; Sari, Piran, & Halpern 1999). Such evidence is rapidly growing up in the recent years (e.g., Bloom, Frail, & Kulkarni 2003 and the references therein).

The structure of GRB jet is currently under heavy debate. Uniform jet model (e.g., Rhoads 1999; Frail et al. 2001) and universal structured jet (USJ) model (e.g., Mészáros, Rees & Wijers 1998; Dai & Gou 2001; Rossi, Lazzati, & Rees 2002; Zhang & Mészáros 2002) are two currently competing models. In the framework of the uniform jet model, jet opening angle is assumed to be different from burst to burst, but energy distribution within a jet is uniform. In the scenario of the structured jet model, it is assumed that the energy (and/or bulk Lorentz factor) distribution within a jet is a function of the angle measured from jet axis. The USJ models suggest that all GRB jets have the same geometric structure and the same energy distribution within jets. In this model, the observational diversity of GRB population is resulted only from differ-

ent viewing angles. Since there are significant dispersions in GRB data, quasi-universal structured jet models are also proposed (Zhang et al. 2004a; Lloyd-Ronning, Dai, & Zhang 2004; and Dai & Zhang 2005). Numerical simulations of jet propagation within collapsar context by Zhang, Woosley, & Heger (2004b) showed that the jet structure seems to be quite complicated, especially when the part of $\theta > 0.1$ rad is taken into account. Such a jet structure is difficult to model by a single component. A jet structure with two or more components is likely to be more real. The observations of GRB 030329 seem to favor such a jet model (Berger et al. 2003). The afterglow lights of this burst show two temporal breaks—one occurs at 0.5 day after the burst trigger in optical afterglow light curve (Price et al. 2003) and the other happens at 9.5 days in the radio afterglow light curve (Berger et al. 2003). Berger et al. (2003) argued that the interpretation for the two breaks requires a two-component jet with $\theta_{c,1} = 0.1$ rad and $\theta_{c,2} = 0.3$ rad. Millimeter observations of this burst also support this jet model (Sheth et al. 2003). Liang & Dai (2004) explained the plausible bimodal distribution of the observed peak energy of νF_ν spectra (E_p) with the similar jet structure model. Huang et al. (2004) and Peng, Königl, & Granot (2004) investigated optical afterglows from such a jet model.

Afterglow observations present more detailed information, but in the late afterglow phase jet structure signatures may be washed out. Combination of prompt gamma-ray emissions and afterglow emissions (especially the early

afterglow emissions) may be a powerful way to test different jet models. Zhang et al. (2004), Lloyd-Ronning et al. (2004), and Dai & Zhang (2005) showed that the observed $E_{\text{iso}} - E_p$ relation (Amati et al. 2002; Sakamoto et al. 2004; Lamb et al. 2005; Liang, Dai & Wu 2004), θ distribution, GRB distributions in (θ, z) -plane and in (E_p, S_γ) -plane, luminosity function, and $\log N - \log P$ distribution are roughly consistent with simulation results based on a quasi-universal Gaussian-like jet model, where E_{iso} is the equivalent-isotropic energy of GRBs, S_γ is the observed fluence in gamma-ray band, and z is the redshift of GRBs. Firmani et al. (2004) constrained the isotropic luminosity function and formation rate of long GRBs by fitting models jointly to both the observed differential peak flux and redshift distributions, and found evidence supporting a jet model intermediate between universal power-law jet model and quasi-universal Gaussian structured model. For the uniform jet model their result is compatible with an angle distribution between 2° and 15° . The USJ model has a power to predict the distribution of viewing angles, $P^{\text{th}}(\theta)$. Perna, Sari, & Frail (2003) utilized this power to test the USJ models, and found that $P^{\text{th}}(\theta)$ is roughly consistent with the observed one derived from a sample of 16 events with θ known from Bloom et al. (2003). However, Nakar, Granot & Guetta (2004) made a further analysis, and found that $P^{\text{th}}(\theta, z)$ does not agree with the observed result. Liang, Wu & Dai (2004) tested the USJ model by simulations based on the $E_{\text{iso}} - E_p$ relation and the assumption of a standard energy budget in GRB jets (Frail et al. 2001). In their simulations, they adopted a high threshold of gamma-ray fluence, and found that simulated $P(\theta)$ and $P(\theta, z)$ are consistent with USJ model predictions.

The largest GRB sample available so far is the BATSE GRB sample. Statistical test with this sample might be more reasonable and reliable. More recently, Ghirlanda, Ghisellini, & Lazzati (2004) discovered a very tight correlation between the gamma-ray energy in GRB jet (E_γ) and E'_p , where $E'_p = E_p \times (1+z)$ (hereafter GGL-relation). It gives an empirical way to estimate the θ of a burst once the E'_p of the bursts is available. In this analysis, we use this relationship to derive the empirical θ distribution of the long BATSE GRB sample, $P^{\text{em}}(\theta)$, and the distribution of these GRBs in the (θ, z) -plane, $P^{\text{em}}(\theta, z)$. We test different USJ models by comparing the model predictions, $P^{\text{th}}(\theta)$ and $P^{\text{th}}(\theta, z)$, with our empirical results. The differences of this analysis from both Perna et al. (2003) and Nakar et al. (2004) are summarized as follows. (1) The sample we used is the long BATSE GRB ($T_{90} > 2$ seconds) sample (1213 bursts) but not the current θ -known GRB sample. The current θ -known GRB sample is too small and it suffers greatly observational biases and sample selection effects (e.g., Broom & Leob 2002; Bloom 2003). (2) The observed GRB rate is taken as a corrected star formation rate proposed by Bloom (2003). When we compute the θ for each burst, we assign a redshift from the observed GRB rate model by a simple simulation. This GRB rate model is also used to calculate the $P(\theta)$ and $P(\theta, z)$ predicted by the USJ models. (3) The θ of each burst is calculated by the GGL-relation. Since the E_p values of most of the bursts in our sample are not available, we derive the E_p values from a relationship between E_p

and hardness ratios (HR). (4) To ensure that the sample should be regarded as a complete one and to make the comparisons between empirical results and model predictions reasonable, we use an “effective” threshold according to our sample selection criteria. (5) Various USJ models have been tested.

This paper is arranged as follows. In section 2, we describe our empirical approach to derive $P^{\text{em}}(\theta)$ and $P^{\text{em}}(\theta, z)$ for the long BATSE GRBs. In section 3, we present the theoretical models of $P^{\text{th}}(\theta)$ and $P^{\text{th}}(\theta, z)$ predicted by the USJ models. Results are presented in section 4, and discussion and conclusions are presented in section 5. Throughout this paper we adopt $\Omega_M = 0.3$, $\Omega_\Lambda = 0.7$, and $H_0 = 71 \text{ km Mpc}^{-1} \text{ s}^{-1}$.

2. EMPIRICAL APPROACH

The energy release in the gamma-ray band of a GRB jet is

$$E_\gamma = E_{\text{iso}}(1 - \cos \theta), \quad (1)$$

and E_{iso} is calculated by

$$E_{\text{iso}} = \frac{4\pi D_L^2 k S_\gamma}{1 + z}, \quad (2)$$

where S_γ is the observed gamma-ray fluence, D_L is the luminosity distance at redshift z , and k is a factor to correct the observed fluence in an instrument band to a standard bandpass in the rest-frame ($1 - 10^4 \text{ keV}$ in this analysis; Bloom et al. 2001). The GGL-relation is,

$$E_{\gamma,50} \simeq [E_{p,2}(1+z)]^{1.5}, \quad (3)$$

where $E_{p,2} = E_p/100 \text{ keV}$ and $E_{\gamma,50} = E_\gamma/10^{50} \text{ ergs}$ (see also, Dai, Liang, & Xu 2004). Although the physics behind this relationship is still poorly understood, the very small dispersion of this relationship makes it a reliable way to estimate the value of θ , once the E_p and z of a burst are well measured. We use this relationship to calculate the θ values of the long BATSE GRBs. Combining Eqs. 1-3, we have

$$\theta = \arccos[1 - \frac{1}{4\pi} \frac{E_{p,2}^{1.5}(1+z)^{2.5}}{D_{L,28}^2 S_{\gamma,-6k}}]. \quad (4)$$

The E_p values of the bursts in our sample are only available for some bright GRBs (Band et al. 1993; Ford et al. 1995; Preece et al. 1998, 2000). It is well known that the distribution of E_p and HR of long BATSE GRBs are narrowly clustered (e.g., Preece et al. 2000). In a previous work we showed that the E_p and HR of those GRBs with a moderate E_p ($100 \sim 1000 \text{ keV}$) and HR ($1.6 \sim 6$, calculated by the fluence measured in the energy band $110\text{-}300 \text{ keV}$ to that in $55\text{-}110 \text{ keV}$) are strongly correlated (Cui, Liang, & Lu 2005). Thus, we estimate the E_p values by the correlation between E_p and HR . We show this correlation for a bright GRB sample of 149 bursts presented by Lloyd-Ronning & Ramirez-Ruiz (2002) in the upper panel of Figure 1. We perform a linear least square fit at 1σ confidence level to the two quantities. We have

$$\log E_p = (1.86 \pm 0.14) + (1.16 \pm 0.09) \log HR \quad (5)$$

with a linear correlation coefficient of 0.74 and a chance probability of $p < 10^{-4}$. The reduced χ^2 is 3.84. We mark

the 1σ region in Figure 1 (gray band). The distribution of the HR for the long BATSE GRB sample is also shown in the lower panel of Figure 1. From Figure 1 we find that most of the GRBs in our sample well follow this relationship.

The redshifts of most GRBs in our sample are unknown. Please note that our purpose is to examine whether or not $P^{\text{em}}(\theta)$ and $P^{\text{em}}(\theta, z)$ are consistent with those predicted by the USJ models. A GRB rate model has to be used in our calculations. To compare the model predictions with the empirical results, the same GRB rate model should be used to derive the empirical results and model results. We assume that the observed GRB rate as a function of redshift for the long GRBs is the same as that derived from the currently redshift-known GRB sample. Bloom (2003) considered a correction factor of D_L^{-2} for $z > 1$ and showed that the models of star formation rate as a function of redshift, SF1, SF2, and SF3, from Porciani & Madau (2001), are consistent with the observed GRB rate. We hence use the same GRB rate model suggested by Bloom (2003) in our calculations. Since the largest redshift of GRBs observed so far is 4.5, we also limit $z \leq 4.5$. For a given burst, its redshift is assigned by a simple Monte Carlo simulation. The procedure of our simulation is as follows: (1) obtain the differential probability of the observed redshift distribution with a bin size of 0.01, $dQ(z)/dz$; (2) derive the accumulated probability distribution of redshift, $Q(z)$ ($0 < Q(z) \leq 1$); (3) generate a random number, m ($0 < m \leq 1$); and (4) assign a z value to the bursts by $z = (z_{i+1} + z_i)/2$, where $Q(z_i) < m$ and $Q(z_{i+1}) > m$. Please note that the redshift assigned by this way for a burst can be used only for a statistical purpose. Such a simulation ensures that $P(\theta)$ and $P(\theta, z)$ predicted by the USJ models and derived from BASTE observations are based on the same GRB rate model.

Based on the analysis above, we calculate the θ values by Eq. 4. The uncertainty of θ is

$$\sigma_\theta = \sqrt{\left(\frac{\partial\theta}{\partial E_p}\right)^2 \sigma_{E_p}^2 + \left(\frac{\partial\theta}{\partial E_{\text{iso}}}\right)^2 \sigma_{E_{\text{iso}}}^2} = \frac{y}{\sqrt{1-x^2}} \sqrt{\left(1.5 \frac{\sigma_{E_p}}{E_p}\right)^2 + \left(\frac{\sigma_{E_{\text{iso}}}}{E_{\text{iso}}}\right)^2} \quad (6)$$

where $y = [E_{p,2}(1+z)]^{1.5}/E_{\text{iso},50}$, $x = 1 - y$. The uncertainties of E_{iso} and E_p are given by

$$\sigma_{E_{\text{iso}}} = \frac{4\pi D_L^2}{1+z} k \sigma_{S_\gamma} \quad (7)$$

and

$$\sigma_{E_p} = \sqrt{\left(\frac{\partial E_p}{\partial a}\right)^2 \sigma_a^2 + \left(\frac{\partial E_p}{\partial b}\right)^2 \sigma_b^2 + \left(\frac{\partial E_p}{\partial HR}\right)^2 \sigma_{HR}^2} = E_p \ln 10 \sqrt{\sigma_a^2 + (\log HR)^2 \sigma_b^2 + \left(\frac{b \sigma_{HR}}{HR \ln 10}\right)^2} \quad (8)$$

respectively, where $a = 1.86 \pm 0.14$ and $b = 1.16 \pm 0.09$. Please note when we calculate the $\sigma_{E_{\text{iso}}}$, the uncertainty of k is ignored since the spectral parameters of the bursts are not available (we assume $\alpha = -1$ and $\beta = -2.3$ to compute k values for all the bursts).

3. THEORETICAL MODELS

The $P(\theta)$ and $P(\theta, z)$ predicted by the USJ models for a given detection threshold are

$$P^{\text{th}}(\theta) = \sin \theta \int_0^{z_{\text{max}}} dz \frac{R_{\text{GRB}}}{1+z} \frac{dV}{dz} \quad (9)$$

and

$$P^{\text{th}}(\theta, z) = \sin \theta \frac{R_{\text{GRB}}}{1+z} \frac{dV}{dz}, \quad (10)$$

respectively, where R_{GRB} is the GRB rate per unit comoving time per unit comoving volume, dV/dz is the comoving volume element at z , and z_{max} is the maximum redshift up to which a burst with viewing angle θ can satisfy the detection threshold. In this analysis, the R_{GRB} is taken as the observed GRB rate suggested by Bloom (2003). The z_{max} is determined by instrument sensitivity and jet structure model. It can be derived from

$$E_\gamma = \int_0^{\pi/2} 4\pi \epsilon(\theta) \sin \theta d\theta = \frac{4\pi D_L^2(z_{\text{max}}) \tilde{S}_{th} k}{1+z_{\text{max}}} (1 - \cos \theta), \quad (11)$$

where $\tilde{S}_{th}$ is the “effective” threshold of the long GRB sample observed by BATSE, and $\epsilon(\theta)$ is energy density in per solid angle as a function of θ . Please note that $\tilde{S}_{th}$ does not correspond to the BATSE threshold. An observed GRB sample is significantly affected by observational biases and sample selection effects, especially when the completeness at low fluxes is considered. The “effective” threshold should correspond to a given sample selection criterion to ensure that the sample should be regarded as a complete one in this threshold. Such a sample should be selected from a sensitive all-sky survey. The fluence distribution of our sample is shown in Figure 2. It is found 95% of the bursts satisfy $S > 3.2 \times 10^{-7}$ ergs. cm^{-2} . We thus take $\tilde{S}_{th}$ as 3.2×10^{-7} ergs. cm^{-2} . The energy density profile $\epsilon(\theta)$ for a single power-law jet, Gaussian jet, and two-components jet are written as follows.

Power-law jet:

$$\epsilon(\theta) = \epsilon_c \left(\frac{\theta}{\theta_c}\right)^{-2} \quad (12)$$

Gaussian jet:

$$\epsilon(\theta) = \epsilon_0 e^{-\theta^2/2\theta_0^2} \quad (13)$$

where θ_0 is a characteristic width of the jet, and ϵ_0 is the maximum value of the energy density.

Two-component jet:

$$\epsilon(\theta) = \epsilon_0 (e^{-\theta^2/2\theta_{c,1}^2} + \lambda e^{-\theta^2/2\theta_{c,2}^2}) \quad (14)$$

where $\theta_{c,1}$ and $\theta_{c,2}$ are respectively characteristic widths of the two components, and λ is the ratio of energy densities in the two components.

Based on empirical approach and theoretical models discussed above, we calculate the empirical and theoretical θ distributions and the distributions of these GRBs in the (θ, z) -plane. By comparing these empirical results with model's predictions, we test these jet models.

Zhang et al. (2004) showed that the current GRB/XRF prompt emission/afterglow data can be described by a quasi-Gaussian type structured jet (or similar structure jet) with a typical opening angle of 0.1 rad and with a

standard jet energy of $\sim 10^{51}$ ergs. In addition, Liang (2004) showed that the fluence of those GRBs with $\theta < 0.1$ rad is almost a constant. We thus take $\theta_0 = 0.1$ rad and $\epsilon_0 = 10^{51}$ ergs for the Gaussian jet. For the power-law jet model, we take $\theta_c = 0.1$ rad and let ϵ_c be variable. We find that $\epsilon_c = 2.3 \times 10^{50}$ ergs yields the best consistency between $P^{\text{em}}(\theta)$ and $P^{\text{th}}(\theta)$. Berger et al. (2003) argued that the observations of GRB 030329 require a two-component explosion with $\theta_{c,1} = 0.1$ and $\theta_{c,2} = 0.3$. Liang & Dai (2004) obtained the similar results based on the E_p distribution observed by BATSE and HETE-2. We thus take $\theta_{c,1} = 0.1$ and $\theta_{c,2} = 0.3$ for the two-component model. The ratio of energy densities in the two components, λ , is suggested to be $10^{-1.7}$ in Liang & Dai (2004). In this analysis we let it be variable in the range of (0.01, 0.1) to derive a good consistency between $P^{\text{em}}(\theta)$ and $P^{\text{th}}(\theta)$. We use different values of λ , and find that $\lambda \sim 0.08$ gives the best consistency. The ϵ_0 in the two-component jet model is taken as $(10 \pm 4) \times 10^{50}$ ergs, where the error is assumed to be normal distributed. These model parameters are summarized in Table 1.

The empirical results and their comparisons to model's predictions are shown in Figs. (3)-(5). The comparisons between $P^{\text{em}}(\theta)$ and $P^{\text{th}}(\theta)$ are shown in the left panels of the three figures, and comparisons between $P^{\text{em}}(\theta, z)$ and $P^{\text{em}}(\theta, z)$ are shown in the right panels. The error bars of $P^{\text{em}}(\theta)$ are calculated by the following method: we first derive $P^{\text{em}}(\theta + \sigma_\theta)$ and $P^{\text{em}}(\theta - \sigma_\theta)$, and then obtain $\sigma_{P^{\text{em}}(\theta)}$ by $\sigma_{P^{\text{em}}(\theta)} = |P^{\text{em}}(\theta)(\theta + \sigma_\theta) - P^{\text{em}}(\theta)(\theta - \sigma_\theta)|/2$. Those GRBs with θ -known from Bloom et al. (2003) are also marked by star symbols in the (θ, z) -planes. The distribution of these GRBs are roughly consistent with the 2σ region of the $P^{\text{em}}(\theta, z)$. We quantify the difference between $P^{\text{em}}(\theta)$ and $P^{\text{th}}(\theta)$ by a Kolmogorov-Smirnov (K-S) test. The result of the K-S test is depicted by a statistic of P_{K-S} : a small value of P_{K-S} indicates a significant difference between the two distributions ($P_{K-S} = 1$ means two distributions are identical, and $P_{K-S} < 1.0 \times 10^{-4}$ suggests that they are significantly different). The K-S test results are also listed in Table 1. For the comparison between $P^{\text{em}}(\theta, z)$ and $P^{\text{em}}(\theta, z)$, we do not have a quantifiable way to estimate the difference but evaluate it by eye instead.

From Fig. (3) and Table 1, we find the empirical $P^{\text{em}}(\theta)$ is quite consistent with $P^{\text{th}}(\theta)$ predicted by the power-law jet model. The K-S test shows $P_{K-S} = 0.575$, strongly indicating an agreement between $P^{\text{em}}(\theta)$ and $P^{\text{th}}(\theta)$. The $P^{\text{em}}(\theta)$ distribution is well fitted by a log-normal function,

$$P^{\text{em}}(\theta) = \frac{0.67}{0.57\sqrt{\pi/2}} e^{-2(\frac{\log \theta + 0.76}{0.57})^2}. \quad (15)$$

However, the $P^{\text{em}}(\theta, z)$ is greatly different from $P^{\text{th}}(\theta, z)$ predicted by this jet model. Even in 1σ regions, they are quite different. The lack of high- z and large- θ (the right-top of the right panel of Fig. 3) make most of this difference. We adjust the parameters of this jet model, but we still do not get an agreement between $P^{\text{em}}(\theta, z)$ and $P^{\text{th}}(\theta, z)$.

Shown in Fig. (4) is the comparisons of the empirical results with those predicted by the Gaussian jet model. It is found that the $P^{\text{th}}(\theta)$ rapidly drops at $\theta > 0.3$ rad. This is significantly different from the empirical result. The K-

S test for the two distributions shows $P_{K-S} = 2.1 \times 10^{-9}$, indicating that the hypothesis of consistency is rejected. Similar to the $P^{\text{th}}(\theta, z)$ predicted by this jet model, the $P^{\text{th}}(\theta, z)$ predicted by the Gaussian jet model is also deviates the $P^{\text{em}}(\theta, z)$. However, comparing the results shown in the right panels of Figures 3 and 4, one can find that the $P^{\text{th}}(\theta, z)$ predicted by the Gaussian jet model is more consistent with $P^{\text{em}}(\theta, z)$ than that predicted by the power-law jet model.

The comparisons of $P^{\text{em}}(\theta)$ and $P^{\text{em}}(\theta, z)$ with those derived from the two-component jet model are shown in Figure 5. The K-S test for the distributions of $P^{\text{em}}(\theta)$ and $P^{\text{th}}(\theta)$ obtains 3.55×10^{-4} , indicating the consistence between the two distributions is marginally accepted. The $P^{\text{th}}(\theta, z)$ predicted by this jet model is also roughly consistent with $P^{\text{em}}(\theta, z)$ in 1σ region.

5. CONCLUSIONS AND DISCUSSION

Based on the assumption that the observed GRB rate is proportional to a corrected star formation rate, we have derived the empirical distributions of long BATSE GRBs, $P^{\text{em}}(\theta)$ and $P^{\text{em}}(\theta, z)$, by the GGL-relation. Our results show that $P^{\text{em}}(\theta)$ is well fitted by a log-normal distribution centering at $\log \theta = -0.76$ with a width of 0.57. We test different USJ models by comparing model predictions, $P^{\text{th}}(\theta)$ and $P^{\text{th}}(\theta, z)$, with the empirical results. To make the comparison reasonable, an “effective” threshold, which corresponds to the sample selection criteria of the long GRB sample, is used. We find that $P^{\text{th}}(\theta)$ predicted by the power-law jet model is well consistent with $P^{\text{em}}(\theta)$, but $P^{\text{th}}(\theta, z)$ predicted by this model is significantly different from $P^{\text{em}}(\theta, z)$. Inversely, the $P^{\text{th}}(\theta, z)$ predicted by a single Gaussian jet model is more consistent with $P^{\text{em}}(\theta, z)$ than that predicted by the power-law jet model, but the $P^{\text{th}}(\theta)$ predicted by this model rapidly drops at $\theta > 0.3$ rad, which greatly deviates $P^{\text{em}}(\theta)$. Both $P^{\text{th}}(\theta)$ and $P^{\text{th}}(\theta, z)$ predicted by a jet model with two Gaussian components are roughly consistent with $P^{\text{em}}(\theta)$ and $P^{\text{em}}(\theta, z)$.

The structure of jet is crucial to understanding the nature of GRBs, such as their rate, luminosity function, and explosion energy. Numerical simulations of jet propagation within collapsar context by Zhang, Woosley, & Heger (2004b) showed that the jet structure seems to be quite complicated. A universal, single-component jet seems to be difficult to describe such a jet structure, and a jet with two (or more) components might be more reasonable. The observations of GRB 030329 (Price et al. 2003; Berger et al. 2003; Sheth et al. 2003) and the bimodal distribution of the observed E_p of HETE-2 bursts (Liang & Dai 2004) are evidence for this jet structure. The results of this work also seem to support this jet model. Zhang et al. (2004a), Lloyd-Ronning et al. (2004), and Dai & Zhang (2005) showed that quasi-universal Gaussian jet can well interpret the observations of the current well follow-up GRB sample. Please note that this sample is a bright GRB sample. The median of the fluence in this sample is $\sim 2.5 \times 10^{-5}$ erg cm $^{-2}$ (Bloom et al. 2003). It is possible that the emissions of these bursts are dominated by the core component of the jet. Thus, a quasi-universal Gaussian jet might well explain the observations of these bursts.

The true GRB rate as a function of redshift is difficult

to establish from the current GRB sample. It is generally suggested that the GRB rate follows the star formation rate. Various models of the star formation rate are presented in the literature. Whether or not the model of the star formation rate affects our results? We use a model of the observed GRB rate suggested by Bloom (2003), who constructed a correction factor of D_L^{-2} when $z > 1$. This factor leads to the observed GRB rate deeply decay when $z > 1$. Thus, different models of the star formation rate may give almost the same observed GRB rate (Bloom 2003), and then the model of the star formation rate does not significantly affect our results. In fact, we focus on the comparison of both results on theoretical bases and on the observational bases. The results of the comparison should not be greatly affected by the model of star formation rate (Liang, Wu & Dai 2004).

Our empirical results are derived by the GGL-relation. This relationship depends on cosmological parameters. In this work we adopt $\Omega_M = 0.3$ and $\Omega_\Lambda = 0.7$. We check if the cosmological parameters significantly affect the results of the comparisons between our empirical results and model predictions. We take $\Omega_M = 0.5$ and $\Omega_\Lambda = 0.5$. In this case, the GGL-relation becomes $E_{\gamma,50} = (0.90 \pm 0.12)E_p^{1.42 \pm 0.10}$ for a GRB sample presented by Xu et al. (2005). We show the comparison between the $P^{\text{em}}(\theta)$ based on this relationship and $P^{\text{th}}(\theta)$ predicted by the power-law jet model in the cosmology model with $\Omega_M = 0.5$ and $\Omega_\Lambda = 0.5$ in Figure 6. The K-S test for the two distributions shows $P_{K-S} = 0.123$, confidently sug-

gesting that they are consistent. Comparing the results shown in Figure 6 to that shown in the left panel of Figure 3, one can find that cosmological parameters do not significantly affect our results.

We should clarify that our empirical approach and theoretical model are independent without tautology. The model predictions are statistical distributions, while the empirical results are based on the relationships related to the spectral properties and energy release of GRBs. The empirical approach and theoretical model are intrinsically different.

Our empirical results are roughly consistent with the results from currently θ -known GRB sample. This is a self-consistent result because the GGL-relation was discovered by this GRB sample. For the bursts in this sample their peak energies, temporal breaks of optical afterglow light curves, and redshifts are well measured. Such a sample must suffers greatly observational biases and sample selection effects, especially when the completeness at low fluxes and the bias of redshift measurement are considered. Band & Preece (2005) argued that the GGL-relation may be an artifact of the selection effects, and these selection effects may favor sub-populations of GRBs. If it is really the case, the selection effects should affect our empirical results.

We thank the anonymous referee for his/her valuable suggestions and comments. We also thank Bing Zhang, Zigao Dai, and Yiping Qin for their helpful discussions. This work was supported by the National Natural Science Foundation of China (Grants 10463001).

REFERENCES

- Amati, L., et al. 2002, *A&A*, 390, 81
 Band, D. L., et al. 1993, *ApJ*, 413, 281
 Band, D. L. & Preece, R. D. 2005, 627, *ApJ*, in press (astro-ph/0501559)
 Berger, E., et al. 2003, *Nature*, 426, 154
 Bloom, J. S. 2003, *AJ*, 125, 2865
 Bloom, J. S., Frail, D. A., & Kulkarni, S. R. 2003, *ApJ*, 594, 674
 Bloom, J. S., Frail, D. A., & Sari, R. 2001, *AJ*, 121, 2879
 Broom, V. & Loeb, A. 2002, *ApJ*, 575, 111
 Cheng, K. S. & Lu, T. 2001, *Chin. J. Astron. Astrophys.* 1, 1
 Cui, X. H., Liang, E. W., & Lu, R. J. 2005, *Chin. J. Astron. Astrophys.*, 2, 151
 Dai, X. Y. & Zhang, B. 2005, *ApJ*, 621, 875
 Dai, Z. G. & Gou, L. J. 2001, *ApJ*, 552, 72
 Dai, Z. G., Liang, E. W., & Xu, D. 2004, *ApJL*, 612, 101
 Firmani, C., Avila-Reese, V., Ghisellini, G., & Tutukov, A. V. 2004, *ApJ*, 611, 1033
 Fishman, G. & Meegan, C. 1995, *ARA&A*, 33, 415
 Ford, L. A., et al. 1995, *ApJ*, 439, 307
 Frail, D. A., et al. 2001, *ApJ*, 562, L55
 Ghirlanda, G., Ghisellini, G., & Lazzati, D. 2004, *ApJ*, 616, 331
 Huang, Y. F., Wu, X. F., Dai, Z. G., Ma, H. T., & Lu, T. 2004, *ApJ*, 605, 300
 Lamb, D. Q., Donaghy, T. Q., & Graziani, C. 2005, *ApJ*, 620, 355
 Liang, E. W. 2004, *MNRAS*, 348, 153
 Liang, E. W. & Dai, Z. G. 2004, *ApJ*, 608, L9
 Liang, E. W., Dai, Z. G., & Wu, X. F. 2004, *ApJ*, 606, L29
 Liang, E. W., Wu, X. F., & Dai, Z. G. 2004, *MNRAS*, 354, 81
 Lloyd-Ronning, N. M. & Ramirez-Ruiz, E. 2002, *ApJ*, 576, 101
 Lloyd-Ronning, N. M., Dai, X. Y., & Zhang, B. 2004, *ApJ*, 601, 371
 Mészáros, P., Rees, M. J., & Wijers, R. A. M. J. 1998, *ApJ*, 499, 301
 Mészáros, P. 2002, *ARA&A*, 40, 137
 Nakar, E., Granot, J., & Guetta, D. 2004, *ApJ*, 606, 37
 Peng, F., Königl, A., & Granot, J. 2004, astro-ph/0410384
 Perna, R., Sari, R., & Frail, D. 2003, *ApJ*, 594, 379
 Piran, T. 1999, *Phys. Rep.* 314, 575
 Piran, T. 2005, *Reviews of Modern Physics*, 76, 1143
 Porciani, C. & Madau, P. 2001, *ApJ*, 548, 522
 Preece, R. D., et al. 1998, *ApJ*, 496, 849
 Preece, R. D., et al. 2000, *ApJS*, 126, 19
 Price, P. A., et al. 2003, *Nature*, 423, 844
 Rhoads, J. E. 1999, *A&AS*, 138, 539
 Rossi, E., Lazzati, D., & Rees, M. J. 2002, *MNRAS*, 332, 945
 Sakamoto, T., et al. 2004, *ApJ*, 602, 875
 Sari, R., Piran, T., & Halpern, J. P. 1999, *ApJ*, 519, L17
 Sheth, K., et al. 2003, *ApJ*, 595, L33
 van Paradijs, J., Kouveliotou, C., & Wijers, R. A. M. J. 2000, *ARA&A*, 38, 379
 Xu, D., Dai, Z. G., & Liang, E. W. 2005, *ApJ*, submitted, astro-ph/0501458
 Zhang, B., & Mészáros, P. 2002, *ApJ*, 571, 876
 Zhang, B., & Mészáros, P. 2004, *International Journal of Modern Physics A*, 19, 2385
 Zhang, B., Dai, X. Y., Lloyd-Ronning, N. M., & Mészáros, P. 2004a, *ApJ*, 601, L119
 Zhang, W., Woosley, S. E., & Heger, A. 2004b, *ApJ*, 608, 365




FIG. 1.— *Upper*: $\log E_p$ as a function of $\log HR$ for a sample of 149 bright GRBs presented by Llyod-Ronning & Ramirez-Ruiz (2002). The solid line is the best fit line, and the grey band marks the 1σ region of the best fit. *Lower*: The number distribution of $\log HR$ for 1213 long BATSE GRBs. Two vertical-dotted lines mark a region within which bursts follow the $E_p - HR$ relationship shown in the upper panel.




FIG. 2.— Number distribution of the observed fluence in energy band 25 – 2000 keV for 1213 long BATSE GRBs.




FIG. 3.— Comparisons between the GRB probability distributions, $P(\theta)$ (left panel) and $P(\theta, z)$ (right panel), obtained by our empirical approach and by the power-law jet model. In the *left* panel, the step line with error bars is our empirical results, and the solid line is the result of the model. The line contours in the *right* panel are our empirical $P(\theta, z)$ (1σ and 2σ) and the grey contours are the results of the jet model. The stars in the *right* panel are the distribution of the bursts with θ being derived by jet break times (from Bloom et al. 2003).




FIG. 4.— Comparisons between the GRB probability distributions, $P(\theta)$ (left panel) and $P(\theta, z)$ (right panel), obtained by our empirical approach and by the Gaussian jet model. In the *left* panel, the step line with error bars is our empirical results, and the solid line is the result of the model. The line contours in the *right* panel are our empirical $P(\theta, z)$ (1σ and 2σ) and the grey contours are the results of the jet model. The stars in the *right* panel are the distribution of the bursts with θ being derived by jet break times (from Bloom et al. 2003).




FIG. 5.— Comparisons between the GRB probability distributions, $P(\theta)$ (left panel) and $P(\theta, z)$ (right panel), obtained by our empirical approach and by the two-component jet model. In the *left* panel, the step line with error bars is our empirical results, and the solid line is the result of the model. The line contours in the *right* panel are our empirical $P(\theta, z)$ (1 σ and 2 σ) and the grey contours are the results of the jet model. The stars in the *right* panel are the distribution of the bursts with θ being derived by jet break times (from Bloom et al. 2003).




FIG. 6.— Comparisons between the θ distributions derived by our empirical approach (step line with error bars) and by the power-law jet model (solid line) in a cosmology with parameters $\Omega_M = 0.5$ and $\Omega_\Lambda = 0.5$.